

EXEMPLAR SOLUTIONS MATHS

Chapter 3: Matrices

Class
12



Powered by



Chapter 3: Matrices

Exercise 3.3

Short Answer (S.A.)

1. If a matrix has 28 elements, what are the possible orders it can have? What if it has 13 elements?

Solution:

For a given matrix of order $m \times n$, it has mn elements, where m and n are natural numbers.

Here we have, $m \times n = 28$

$(m, n) = \{(1, 28), (2, 14), (4, 7), (7, 4), (14, 2), (28, 1)\}$

So, the possible orders are $1 \times 28, 2 \times 14, 4 \times 7, 7 \times 4, 14 \times 2, 28 \times 1$.

Also, if it has 13 elements, then $m \times n = 13$

$(m, n) = \{(1, 13), (13, 1)\}$

Thus, the possible orders are $1 \times 13, 13 \times 1$.

2. In the matrix $A = \begin{bmatrix} a & 1 & x \\ 2 & \sqrt{3} & x^2 - y \\ 0 & 5 & \frac{-2}{5} \end{bmatrix}$, write :

(i) The order of the matrix A

(ii) The number of elements

(iii) Write elements a_{23}, a_{31}, a_{12}

Solution:

For the given matrix,

(i) The order of the matrix A is 3×3 .

(ii) The number of elements of the matrix $= 3 \times 3 = 9$

(iii) Elements: $a_{23} = x^2 - y, a_{31} = 0, a_{12} = 1$

3. Construct $a_{2 \times 2}$ matrix where

(i) $a_{ij} = (i - 2j)^2 / 2$

(ii) $a_{ij} = |-2i + 3j|$

Solution:

We have,

$$A = [a_{ij}]_{2 \times 2}$$

(i) Such that, $a_{ij} = (i - 2j)^2 / 2$; where $1 \leq i \leq 2; 1 \leq j \leq 2$

So, the terms of the matrix are

$$a_{11} = \frac{(1-2)^2}{2} = \frac{1}{2} \quad a_{12} = \frac{(1-2 \times 2)^2}{2} = \frac{9}{2}$$

$$a_{21} = \frac{(2-2 \times 1)^2}{2} = 0 \quad a_{22} = \frac{(2-2 \times 2)^2}{2} = 2$$

Therefore, $A = \begin{bmatrix} \frac{1}{2} & \frac{9}{2} \\ 0 & 2 \end{bmatrix}$

(ii) Here, $a_{ij} = |-2i + 3j|$

So, the terms of the matrix are

$$a_{11} = |-2 \times 1 + 3 \times 1| = 1 \quad a_{12} = |-2 \times 1 + 3 \times 2| = 4$$

$$a_{21} = |-2 \times 2 + 3 \times 1| = 1 \quad a_{22} = |-2 \times 2 + 3 \times 2| = 2$$

Therefore, $A = \begin{bmatrix} 1 & 4 \\ 1 & 2 \end{bmatrix}$

4. Construct a 3×2 matrix whose elements are given by $a_{ij} = e^{i \cdot x} \sin jx$

Solution:

Let A be a 3×2 matrix

Such that, $a_{ij} = e^{i \cdot x} \sin jx$; where where $1 \leq i \leq 3$; $1 \leq j \leq 2$

So, the terms are given as

$$a_{11} = e^x \sin x \quad a_{12} = e^x \sin 2x$$

$$a_{21} = e^{2x} \sin x \quad a_{22} = e^{2x} \sin 2x$$

$$a_{31} = e^{3x} \sin x \quad a_{32} = e^{3x} \sin 2x$$

Therefore, $A = \begin{bmatrix} e^x \sin x & e^x \sin 2x \\ e^{2x} \sin x & e^{2x} \sin 2x \\ e^{3x} \sin x & e^{3x} \sin 2x \end{bmatrix}$

5. Find values of a and b if $A = B$, where

$$A = \begin{bmatrix} a+4 & 3b \\ 8 & -6 \end{bmatrix}, \quad B = \begin{bmatrix} 2a+2 & b^2+2 \\ 8 & b^2-5b \end{bmatrix}$$

Solution:

Given, matrix A = matrix B

Then their corresponding elements are equal.

So, we have

$$a_{11} = b_{11}; a + 4 = 2a + 2 \Rightarrow a = 2$$

$$a_{12} = b_{12}; 3b = b^2 + 2 \Rightarrow b^2 - 3b + 2 = 0 \Rightarrow b = 1, 2$$

$$a_{22} = b_{22}; -6 = b^2 - 5b \Rightarrow b^2 - 5b + 6 = 0 \Rightarrow b = 2, 3$$

Hence, $a = 2$ and $b = 2$ (common value)

6. If possible, find the sum of the matrices A and B, where $A = \begin{bmatrix} \sqrt{3} & 1 \\ 2 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} x & y & z \\ a & b & 6 \end{bmatrix}$

Solution:

The given two matrices A and B are of different orders. Two matrices can be added only if order of both the matrices is same. Thus, the sum of matrices A and B is not possible.

7. If $X = \begin{bmatrix} 3 & 1 & -1 \\ 5 & -2 & -3 \end{bmatrix}$ and $Y = \begin{bmatrix} 2 & 1 & -1 \\ 7 & 2 & 4 \end{bmatrix}$, find

(i) $X + Y$

(ii) $2X - 3Y$

(iii) A matrix Z such that $X + Y + Z$ is a zero matrix.

Solution:

Given, $X = \begin{bmatrix} 3 & 1 & -1 \\ 5 & -2 & -3 \end{bmatrix}_{2 \times 3}$ and $Y = \begin{bmatrix} 2 & 1 & -1 \\ 7 & 2 & 4 \end{bmatrix}_{2 \times 3}$

(i) $X + Y = \begin{bmatrix} 3+2 & 1+1 & -1-1 \\ 5+7 & -2+2 & -3+4 \end{bmatrix} = \begin{bmatrix} 5 & 2 & -2 \\ 12 & 0 & 1 \end{bmatrix}$

(ii) $2X - 3Y = 2 \begin{bmatrix} 3 & 1 & -1 \\ 5 & -2 & -3 \end{bmatrix} - 3 \begin{bmatrix} 2 & 1 & -1 \\ 7 & 2 & 4 \end{bmatrix}$

$$= \begin{bmatrix} 6 & 2 & -2 \\ 10 & -4 & -6 \end{bmatrix} - \begin{bmatrix} 6 & 3 & -3 \\ 21 & 6 & 12 \end{bmatrix}$$

$$= \begin{bmatrix} 6-6 & 2-3 & -2+3 \\ 10-21 & -4-6 & -6-12 \end{bmatrix} = \begin{bmatrix} 0 & -1 & 1 \\ -11 & -10 & -18 \end{bmatrix}$$

(iii) $X + Y = \begin{bmatrix} 5 & 2 & -2 \\ 12 & 0 & 1 \end{bmatrix}$

Also, $X + Y + Z = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

So, Z is the additive inverse of $(X + Y)$ or negative of $(X + Y)$.

Therefore, $Z = -(X + Y) = \begin{bmatrix} -5 & -2 & 2 \\ -12 & 0 & -1 \end{bmatrix}$

8. Find non-zero values of x satisfying the matrix equation:

$$x \begin{bmatrix} 2x & 2 \\ 3 & x \end{bmatrix} + 2 \begin{bmatrix} 8 & 5x \\ 4 & 4x \end{bmatrix} = 2 \begin{bmatrix} (x^2 + 8) & 24 \\ (10) & 6x \end{bmatrix}.$$

Solution:

Given,

$$\begin{aligned} x \begin{bmatrix} 2x & 2 \\ 3 & x \end{bmatrix} + 2 \begin{bmatrix} 8 & 5x \\ 4 & 4x \end{bmatrix} &= 2 \begin{bmatrix} x^2 + 8 & 24 \\ 10 & 6x \end{bmatrix} \\ \Rightarrow \begin{bmatrix} 2x^2 & 2x \\ 3x & x^2 \end{bmatrix} + \begin{bmatrix} 16 & 10x \\ 8 & 8x \end{bmatrix} &= \begin{bmatrix} 2x^2 + 16 & 48 \\ 20 & 12x \end{bmatrix} \\ \Rightarrow \begin{bmatrix} 2x^2 + 16 & 2x + 10x \\ 3x + 8 & x^2 + 8x \end{bmatrix} &= \begin{bmatrix} 2x^2 + 16 & 48 \\ 20 & 12x \end{bmatrix} \end{aligned}$$

On comparing the corresponding elements, we get

$$2x + 10x = 48$$

$$12x = 48$$

$$\text{Thus, } x = 4$$

It's also seen that this value of x also satisfies the equation $3x + 8 = 20$ and $x^2 + 8x = 12x$.

Therefore, $x = 4$ (common) is the solution of the given matrix equation.

9. If $A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, show that $(A + B)(A - B) \neq A^2 - B^2$

Solution:

Given, $A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$

So, $(A + B) = \begin{bmatrix} 0+0 & 1-1 \\ 1+1 & 1+0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 2 & 1 \end{bmatrix}$

And, $(A - B) = \begin{bmatrix} 0-0 & 1+1 \\ 1-1 & 1-0 \end{bmatrix} = \begin{bmatrix} 0 & 2 \\ 0 & 1 \end{bmatrix}$

$$(A + B) \cdot (A - B) = \begin{bmatrix} 0 & 0 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 0 & 2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0+0 & 0+0 \\ 0+0 & 4+1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 5 \end{bmatrix} \quad \dots (i)$$

Also, $A^2 = A \cdot A$

$$= \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 0+1 & 0+1 \\ 0+1 & 1+1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$$

And, $B^2 = B \cdot B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0-1 & 0+0 \\ 0+0 & -1+0 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$

Therefore, $A^2 - B^2 = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} - \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix} \quad \dots(ii)$

Hence, from (i) and (ii),
 $(A + B)(A - B) \neq A^2 - B^2$

10. Find the value of x if

$$\begin{bmatrix} 1 & x & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 2 \\ 2 & 5 & 1 \\ 15 & 3 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ x \end{bmatrix} = O.$$

Solution:

Given, $\begin{bmatrix} 1 & x & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 2 \\ 2 & 5 & 1 \\ 15 & 3 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ x \end{bmatrix} = O$

$$\Rightarrow [1 + 2x + 15 \quad 3 + 5x + 3 \quad 2 + x + 2] \begin{bmatrix} 1 \\ 2 \\ x \end{bmatrix} = O$$

$$\Rightarrow [16 + 2x \quad 5x + 6 \quad x + 4] \begin{bmatrix} 1 \\ 2 \\ x \end{bmatrix} = O$$

$$[16 + 2x + 10x + 12 + x^2 + 4x] = 0$$

$$[x^2 + 16x + 28] = 0$$

$$x^2 + 16x + 28 = 0$$

$$(x + 2)(x + 14) = 0$$

Therefore, $x = -2, -14$

11. Show that $A = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$ **satisfies the equation** $A^2 - 3A - 7I = 0$ **and hence find** A^{-1} .

Solution:

Given,

$$A = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$$

$$\text{So, } A^2 = A \cdot A = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} \cdot \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} = \begin{bmatrix} 25-3 & 15-6 \\ -5+2 & -3+4 \end{bmatrix} = \begin{bmatrix} 22 & 9 \\ -3 & 1 \end{bmatrix}$$

$$3A = 3 \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} = \begin{bmatrix} 15 & 9 \\ -3 & -6 \end{bmatrix}$$

$$\text{And, } 7I = 7 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$$

$$\begin{aligned} \text{Hence, } A^2 - 3A - 7I &= \begin{bmatrix} 22 & 9 \\ -3 & 1 \end{bmatrix} - \begin{bmatrix} 15 & 9 \\ -3 & -6 \end{bmatrix} - \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} \\ &= \begin{bmatrix} 22-15-7 & 9-9-0 \\ -3+3-0 & 1+6-7 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = O \end{aligned}$$

Now,

$$A^2 - 3A - 7I = 0$$

Multiplying both sides with A^{-1} , we get

$$A^{-1} [A^2 - 3A - 7I] = A^{-1} \cdot 0$$

$$A^{-1} \cdot A \cdot A - 3A^{-1} \cdot A - 7A^{-1} \cdot I = 0$$

$$I \cdot A - 3I - 7A^{-1} = 0 \quad [\text{As } A^{-1} \cdot A = I]$$

$$A - 3I - 7A^{-1} = 0$$

$$7A^{-1} = A - 3I$$

$$7A^{-1} = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} - \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ -1 & -5 \end{bmatrix}$$

$$\text{Therefore, } A^{-1} = \frac{1}{7} \begin{bmatrix} 2 & 3 \\ -1 & -5 \end{bmatrix}$$

12. Find the matrix A satisfying the matrix equation:

$$\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} A \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Solution:

$$\text{Given, } \begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} A \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\text{Let } P = \begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} \quad Q = \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} \quad \text{and } I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Now,

$$P^{-1}PAQ = P^{-1}I$$

$$\text{So, } IAQ = P^{-1}$$

$$AQ = P^{-1}$$

$$AQQ^{-1} = P^{-1}Q^{-1}$$

$$AI = P^{-1}Q^{-1}$$

$$A = P^{-1}Q^{-1}$$

$$\text{Now adj. } P = \begin{bmatrix} 2 & -1 \\ -3 & 2 \end{bmatrix} \text{ and } |P| = 1$$

$$\text{Hence, } P^{-1} = \begin{bmatrix} 2 & -1 \\ -3 & 2 \end{bmatrix}$$

$$\text{Also adj. } Q = \begin{bmatrix} -3 & -2 \\ -5 & -3 \end{bmatrix} \text{ and } |Q| = -1$$

$$\text{Hence, } Q^{-1} = \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix}$$

$$\begin{aligned} \text{Thus, } A &= P^{-1}Q^{-1} \\ &= \begin{bmatrix} 2 & -1 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix} = \begin{bmatrix} 6-5 & 4-3 \\ -9+10 & -6+6 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \end{aligned}$$

$$13. \text{ Find A, if } \begin{bmatrix} 4 \\ 1 \\ 3 \end{bmatrix} A = \begin{bmatrix} -4 & 8 & 4 \\ -1 & 2 & 1 \\ -3 & 6 & 3 \end{bmatrix}$$

Solution:

$$\text{Given, } \begin{bmatrix} 4 \\ 1 \\ 3 \end{bmatrix} A = \begin{bmatrix} -4 & 8 & 4 \\ -1 & 2 & 1 \\ -3 & 6 & 3 \end{bmatrix}$$

Now, let $A = [x \ y \ z]$

$$\text{So, } \begin{bmatrix} 4 \\ 1 \\ 3 \end{bmatrix} [xyz] = \begin{bmatrix} -4 & 8 & 4 \\ -1 & 2 & 1 \\ -3 & 6 & 3 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 4x & 4y & 4z \\ x & y & z \\ 3x & 3y & 3z \end{bmatrix} = \begin{bmatrix} -4 & 8 & 4 \\ -1 & 2 & 1 \\ -3 & 6 & 3 \end{bmatrix}$$

On comparing elements of both sides, we have

$$4x = -4 \Rightarrow x = -1$$

$$4y = 8 \Rightarrow y = 2$$

$$\text{And, } 4z = 4 \Rightarrow z = 1$$

$$\text{Therefore, } A = \begin{bmatrix} -1 & 2 & 1 \end{bmatrix}$$

14. If $A = \begin{bmatrix} 3 & -4 \\ 1 & 1 \\ 2 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 & 2 \\ 1 & 2 & 4 \end{bmatrix}$, then verify $(BA)^2 \neq B^2 A^2$

Solution:

The given matrices A has order 3×2 and B has order 2×3 .

So, BA is defined and will have order 3×3 .

But, A^2 and B^2 are not defined as the orders don't satisfy the multiplication condition.

Hence, $(BA)^2 \neq B^2 A^2$

15. If possible, find BA and AB where

$$A = \begin{bmatrix} 2 & 1 & 2 \\ 1 & 2 & 4 \end{bmatrix}, B = \begin{bmatrix} 4 & 1 \\ 2 & 3 \\ 1 & 2 \end{bmatrix}$$

Solution:

$$\text{Given, } A = \begin{bmatrix} 2 & 1 & 2 \\ 1 & 2 & 4 \end{bmatrix}_{2 \times 3} \text{ and } B = \begin{bmatrix} 4 & 1 \\ 2 & 3 \\ 1 & 2 \end{bmatrix}_{3 \times 2}$$

So, AB and BA both are defined

$$\text{Now, } AB = \begin{bmatrix} 2 & 1 & 2 \\ 1 & 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & 1 \\ 2 & 3 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 8+2+2 & 2+3+4 \\ 4+4+4 & 1+6+8 \end{bmatrix} = \begin{bmatrix} 12 & 9 \\ 12 & 15 \end{bmatrix}$$

$$\begin{aligned} \text{And, } BA &= \begin{bmatrix} 4 & 1 \\ 2 & 3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 1 & 2 \\ 1 & 2 & 4 \end{bmatrix} \\ &= \begin{bmatrix} 8+1 & 4+2 & 8+4 \\ 4+3 & 2+6 & 4+12 \\ 2+2 & 1+4 & 2+8 \end{bmatrix} = \begin{bmatrix} 9 & 6 & 12 \\ 7 & 8 & 16 \\ 4 & 5 & 10 \end{bmatrix} \end{aligned}$$

16. Show by an example that for $A \neq 0$, $B \neq 0$, $AB = 0$.

Solution:

$$\text{Let } A = \begin{bmatrix} 0 & 1 \\ 0 & 2 \end{bmatrix} \neq O \text{ and } B = \begin{bmatrix} -1 & 1 \\ 0 & 0 \end{bmatrix} \neq O$$

$$\text{So, the product } AB = \begin{bmatrix} 0 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} -1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = O$$

- Hence Proved

17. Given $A = \begin{bmatrix} 2 & 4 & 0 \\ 3 & 9 & 6 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 4 \\ 2 & 8 \\ 1 & 3 \end{bmatrix}$. Is $(AB)' = B' A'$?

Solution:

$$\text{Given, } A = \begin{bmatrix} 2 & 4 & 0 \\ 3 & 9 & 6 \end{bmatrix}_{2 \times 3} \text{ and } B = \begin{bmatrix} 1 & 4 \\ 2 & 8 \\ 1 & 3 \end{bmatrix}_{3 \times 2}$$

So, their product is

$$AB = \begin{bmatrix} 2 & 4 & 0 \\ 3 & 9 & 6 \end{bmatrix} \begin{bmatrix} 1 & 4 \\ 2 & 8 \\ 1 & 3 \end{bmatrix} = \begin{bmatrix} 2+8+0 & 8+32+0 \\ 3+18+6 & 12+72+18 \end{bmatrix} = \begin{bmatrix} 10 & 40 \\ 27 & 102 \end{bmatrix}$$

$$\text{And, } (AB)' = \begin{bmatrix} 10 & 27 \\ 40 & 102 \end{bmatrix} \quad \dots(i)$$

$$\text{Also, } B' = \begin{bmatrix} 1 & 2 & 1 \\ 4 & 8 & 3 \end{bmatrix}_{2 \times 3} \text{ and } A' = \begin{bmatrix} 2 & 3 \\ 4 & 9 \\ 0 & 6 \end{bmatrix}_{3 \times 2}$$

$$\begin{aligned} \text{Therefore, } B'A' &= \begin{bmatrix} 1 & 2 & 1 \\ 4 & 8 & 3 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 4 & 9 \\ 0 & 6 \end{bmatrix} \\ &= \begin{bmatrix} 2+8+0 & 3+18+6 \\ 8+32+0 & 12+72+18 \end{bmatrix} \\ &= \begin{bmatrix} 10 & 27 \\ 40 & 102 \end{bmatrix} \quad \dots(ii) \end{aligned}$$

From (i) and (ii), we have

$$(AB)' = B' A'$$

18. Solve for x and y:

$$x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} 3 \\ 5 \end{bmatrix} + \begin{bmatrix} -8 \\ -11 \end{bmatrix} = O$$

Solution:

$$\text{Given, } x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} 3 \\ 5 \end{bmatrix} + \begin{bmatrix} -8 \\ -11 \end{bmatrix} = O$$

$$\begin{bmatrix} 2x \\ x \end{bmatrix} + \begin{bmatrix} 3y \\ 5y \end{bmatrix} + \begin{bmatrix} -8 \\ -11 \end{bmatrix} = O \quad \text{(Multiplying the variables with the matrices)}$$

$$\text{So, } \begin{bmatrix} 2x + 3y - 8 \\ x + 5y - 11 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \text{(Addition of matrices)}$$

Now, we have

$$2x + 3y - 8 = 0 \dots (1) \text{ and}$$

$$x + 5y - 11 = 0 \dots (2)$$

On solving the equations (1) and (2), we get

$$x = 1 \text{ and } y = 2$$

19. If X and Y are 2 x 2 matrices, then solve the following matrix equations for X and Y

$$2X + 3Y = \begin{bmatrix} 2 & 3 \\ 4 & 0 \end{bmatrix}, 3X + 2Y = \begin{bmatrix} -2 & 2 \\ 1 & -5 \end{bmatrix}$$

Solution:

$$\text{Given, } 2X + 3Y = \begin{bmatrix} 2 & 3 \\ 4 & 0 \end{bmatrix} \dots (i)$$

$$\text{and } 3X + 2Y = \begin{bmatrix} -2 & 2 \\ 1 & -5 \end{bmatrix} \dots (ii)$$

On subtracting equations (i) and (ii), we get

$$(3X + 2Y) - (2X + 3Y) = \begin{bmatrix} -2 - 2 & 2 - 3 \\ 1 - 4 & -5 - 0 \end{bmatrix}$$

$$\text{Thus, } X - Y = \begin{bmatrix} -4 & -1 \\ -3 & -5 \end{bmatrix} \dots (iii)$$

On adding equations (i) and (ii), we get

$$5X + 5Y = \begin{bmatrix} 0 & 5 \\ 5 & -5 \end{bmatrix}$$

$$X + Y = \frac{1}{5} \begin{bmatrix} 0 & 5 \\ 5 & -5 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \dots (iv)$$

On adding equations (iii) and (iv), we get

$$(X - Y) + (X + Y) = \begin{bmatrix} -4 & 0 \\ -2 & -6 \end{bmatrix}$$

$$2X = \begin{bmatrix} -4 & 0 \\ -2 & -6 \end{bmatrix}$$

$$\text{Thus, } X = \begin{bmatrix} -2 & 0 \\ -1 & -3 \end{bmatrix}$$

From equation (iv), we get

$$\begin{bmatrix} -2 & 0 \\ -1 & -3 \end{bmatrix} + Y = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}$$

$$Y = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} - \begin{bmatrix} -2 & 0 \\ -1 & -3 \end{bmatrix}$$

$$\text{Thus, } Y = \begin{bmatrix} 2 & 1 \\ 2 & 2 \end{bmatrix}$$

20. If $A = \begin{bmatrix} 3 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 7 & 3 \end{bmatrix}$, then find a non-zero matrix C such that $AB = AC$.

Solution:

Given, $A = \begin{bmatrix} 3 & 5 \end{bmatrix}_{1 \times 2}$ and $B = \begin{bmatrix} 7 & 3 \end{bmatrix}_{1 \times 2}$

For $AC = BC$

We have order of $C = 2 \times n$

For $n = 1$, let $C = \begin{bmatrix} x \\ y \end{bmatrix}$

$$\therefore AC = \begin{bmatrix} 3 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = [3x + 5y]$$

$$\text{And } BC = \begin{bmatrix} 7 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = [7x + 3y]$$

For $AC = BC$,

$$[3x + 5y] = [7x + 3y]$$

$$3x + 5y = 7x + 3y$$

$$4x = 2y$$

$$x = \frac{1}{2}y$$

$$y = 2x$$

$$\text{Hence, } C = \begin{bmatrix} x \\ 2x \end{bmatrix}$$

It's seen that on taking C of order $2 \times 1, 2 \times 2, 2 \times 3, \dots$, we get

$$C = \begin{bmatrix} x \\ 2x \end{bmatrix}, \begin{bmatrix} x & x \\ 2x & 2x \end{bmatrix}, \begin{bmatrix} x & x & x \\ 2x & 2x & 2x \end{bmatrix} \dots$$

In general,

$$C = \begin{bmatrix} k \\ 2k \end{bmatrix}, \begin{bmatrix} k & k \\ 2k & 2k \end{bmatrix} \text{ etc } \dots$$

21. Give an example of matrices A, B and C such that $AB = AC$, where A is non-zero matrix, but $B \neq C$.

Solution:

Let $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$ $[\because B \neq C]$

$$AB = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \quad \dots(i)$$

and $AC = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \quad \dots(ii)$

From (i) and (ii),

Hence, $AB = AC$ but $B \neq C$.

22. If $A = \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 3 \\ 3 & -4 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix}$,

(i) $(AB)C = A(BC)$

(ii) $A(B + C) = AB + AC$.

Solution:

Given, $A = \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 3 \\ 3 & -4 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix}$

(i) $AB = \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 3 & -4 \end{bmatrix} = \begin{bmatrix} 2+6 & 3-8 \\ -4+3 & -6-4 \end{bmatrix} = \begin{bmatrix} 8 & -5 \\ -1 & -10 \end{bmatrix}$

and $(AB)C = \begin{bmatrix} 8 & -5 \\ -1 & -10 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 8+5 & 0 \\ -1+10 & 0 \end{bmatrix} = \begin{bmatrix} 13 & 0 \\ 9 & 0 \end{bmatrix} \quad \dots(i)$

Again, $(BC) = \begin{bmatrix} 2 & 3 \\ 3 & -4 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 2-3 & 0 \\ 3+4 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 7 & 0 \end{bmatrix}$

And $A(BC) = \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 7 & 0 \end{bmatrix} = \begin{bmatrix} -1+14 & 0 \\ 2+7 & 0 \end{bmatrix} = \begin{bmatrix} 13 & 0 \\ 9 & 0 \end{bmatrix} \quad \dots(ii)$

From (i) and (ii), we get

Hence, $(AB)C = A(BC)$

$$(ii) \quad B + C = \begin{bmatrix} 2 & 3 \\ 3 & -4 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 2 & -4 \end{bmatrix}$$

Now,

$$\begin{aligned} A \cdot (B + C) &= \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 3 & 3 \\ 2 & -4 \end{bmatrix} \\ &= \begin{bmatrix} 3+4 & 3-8 \\ -6+2 & -6-4 \end{bmatrix} \\ &= \begin{bmatrix} 7 & -5 \\ -4 & -10 \end{bmatrix} \quad \dots(iii) \end{aligned}$$

$$AB = \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 3 & -4 \end{bmatrix} = \begin{bmatrix} 2+6 & 3-8 \\ -4+3 & -6-4 \end{bmatrix} = \begin{bmatrix} 8 & -5 \\ -1 & -10 \end{bmatrix}$$

$$\text{and, } AC = \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 1-2 & 0 \\ -2-1 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ -3 & 0 \end{bmatrix}$$

$$\text{Thus, } AB + AC = \begin{bmatrix} 8 & -5 \\ -1 & -10 \end{bmatrix} + \begin{bmatrix} -1 & 0 \\ -3 & 0 \end{bmatrix} = \begin{bmatrix} 7 & -5 \\ -4 & -10 \end{bmatrix} \quad \dots(iv)$$

Hence from equation (iii) and (iv), we have

$$A \cdot (B + C) = A \cdot B + A \cdot C$$

$$23. \quad \text{If } P = \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix} \text{ and } Q = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, \text{ prove that } PQ = \begin{bmatrix} xa & 0 & 0 \\ 0 & yb & 0 \\ 0 & 0 & zc \end{bmatrix} = QP.$$

Solution:

$$\text{Given, } P = \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix} \text{ and } Q = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$$

It's seen that both P and Q are diagonal matrices.

We know that, for diagonal matrices elements of product matrix are obtained by multiplying elements of matrices in the principal diagonal.

Hence,

$$\begin{aligned} PQ &= \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix} \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} \\ &= \begin{bmatrix} xa & 0 & 0 \\ 0 & yb & 0 \\ 0 & 0 & zc \end{bmatrix} = \begin{bmatrix} ax & 0 & 0 \\ 0 & by & 0 \\ 0 & 0 & cz \end{bmatrix} = QP \end{aligned}$$

Therefore, $PQ = QP$

24. If: $[2 \ 1 \ 3] \begin{bmatrix} -1 & 0 & -1 \\ -1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = A$, find A.

Solution:

Given, $[2 \ 1 \ 3]_{1 \times 3} \begin{bmatrix} -1 & 0 & -1 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}_{3 \times 3} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}_{3 \times 1} = A$

So, $[2 \ 1 \ 3]_{1 \times 3} \begin{bmatrix} -1+0+1 \\ -1+0+0 \\ 0+0-1 \end{bmatrix}_{3 \times 1} = A$

$[2 \ 1 \ 3]_{1 \times 3} \begin{bmatrix} 0 \\ -1 \\ -1 \end{bmatrix}_{3 \times 1} = A$

$[0 \ -1 \ -3] = A$

Thus, $A = [-4]$

25. If $A = [2 \ 1]$, $B = \begin{bmatrix} 5 & 3 & 4 \\ 8 & 7 & 6 \end{bmatrix}$ and $C = \begin{bmatrix} -1 & 2 & 1 \\ 1 & 0 & 2 \end{bmatrix}$, verify that

$A(B + C) = (AB + AC)$.

Solution:

Given, $A = [2 \ 1]$, $B = \begin{bmatrix} 5 & 3 & 4 \\ 8 & 7 & 6 \end{bmatrix}$ and $C = \begin{bmatrix} -1 & 2 & 1 \\ 1 & 0 & 2 \end{bmatrix}$

Now,

$A(B + C) = [2 \ 1] \begin{bmatrix} 5-1 & 3+2 & 4+1 \\ 8+1 & 7+0 & 6+2 \end{bmatrix}$

$= [2 \ 1] \begin{bmatrix} 4 & 5 & 5 \\ 9 & 7 & 8 \end{bmatrix}$

$= [8 + 9 \ 10 + 7 \ 10 + 8]$

$= [17 \ 17 \ 18]$

...(i)

And,

$AB = [2 \ 1] \begin{bmatrix} 5 & 3 & 4 \\ 8 & 7 & 6 \end{bmatrix} = [10 + 8 \ 6 + 7 \ 8 + 6] = [18 \ 13 \ 14]$

And,

$$AC = [2 \ 1] \begin{bmatrix} -1 & 2 & 1 \\ 1 & 0 & 2 \end{bmatrix} = [-2 + 1 \ 4 + 0 \ 2 + 2] = [-1 \ 4 \ 4]$$

$$\text{So, } AB + AC = [18 \ 13 \ 14] + [-1 \ 4 \ 4] \\ = [17 \ 17 \ 18]$$

... (ii)

From equations (i) and (ii),

$$A(B + C) = (AB + AC)$$

Hence, verified

26. If $A = \begin{bmatrix} 1 & 0 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 1 \end{bmatrix}$, then verify that $A^2 + A = A(A + I)$, where I is 3×3 unit matrix.

Solution:

$$\text{Given, } A = \begin{bmatrix} 1 & 0 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 1 \end{bmatrix}$$

$$\text{So, } A^2 = A \cdot A$$

$$\text{Thus, } = \begin{bmatrix} 1 & 0 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1+0+0 & 0+0-1 & -1+0-1 \\ 2+2+0 & 0+1+3 & -2+3+3 \\ 0+2+0 & 0+1+1 & 0+3+1 \end{bmatrix} = \begin{bmatrix} 1 & -1 & -2 \\ 4 & 4 & 4 \\ 2 & 2 & 4 \end{bmatrix}$$

$$A^2 + A = \begin{bmatrix} 1 & -1 & -2 \\ 4 & 4 & 4 \\ 2 & 2 & 4 \end{bmatrix} + \begin{bmatrix} 1 & 0 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 2 & -1 & -3 \\ 6 & 5 & 7 \\ 2 & 3 & 5 \end{bmatrix} \quad \dots (i)$$

$$\text{Now, } A + I = \begin{bmatrix} 1 & 0 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 0 & -1 \\ 2 & 2 & 3 \\ 0 & 1 & 2 \end{bmatrix}$$

$$\text{So, } A(A + I) = \begin{bmatrix} 1 & 0 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 & -1 \\ 2 & 2 & 3 \\ 0 & 1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 2+0+0 & 0+0-1 & -1+0-2 \\ 4+2+0 & 0+2+3 & -2+3+6 \\ 0+2+0 & 0+2+1 & 0+3+2 \end{bmatrix} = \begin{bmatrix} 2 & -1 & -3 \\ 6 & 5 & 7 \\ 2 & 3 & 5 \end{bmatrix} \dots(ii)$$

From (i) and (ii), we get

$$A^2 + A = A(A + I)$$

27. If $A = \begin{bmatrix} 0 & -1 & 2 \\ 4 & 3 & -4 \end{bmatrix}$ and $B = \begin{bmatrix} 4 & 0 \\ 1 & 3 \\ 2 & 6 \end{bmatrix}$, then verify that:

(i) $(A')' = A$

(ii) $(AB)' = B'A'$

(iii) $(kA)' = (kA')$

Solution:

Given, $A = \begin{bmatrix} 0 & -1 & 2 \\ 4 & 3 & -4 \end{bmatrix}$ and $B = \begin{bmatrix} 4 & 0 \\ 1 & 3 \\ 2 & 6 \end{bmatrix}$

(i) We have to verify that, $(A')' = A$

So,

$$A' = \begin{bmatrix} 0 & 4 \\ -1 & 3 \\ 2 & -4 \end{bmatrix}$$

And, $(A')' = \begin{bmatrix} 0 & -1 & 2 \\ 4 & 3 & -4 \end{bmatrix} = A$

(ii) We have to verify that, $(AB)' = B'A'$

So,

$$AB = \begin{bmatrix} 0 & -1 & 2 \\ 4 & 3 & -4 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 1 & 3 \\ 2 & 6 \end{bmatrix} = \begin{bmatrix} 3 & 9 \\ 11 & -15 \end{bmatrix}$$

$$(AB)' = \begin{bmatrix} 3 & 11 \\ 9 & -15 \end{bmatrix}$$

and, $B'A' = \begin{bmatrix} 4 & 1 & 2 \\ 0 & 3 & 6 \end{bmatrix} \begin{bmatrix} 0 & 4 \\ -1 & 3 \\ 2 & -4 \end{bmatrix} = \begin{bmatrix} 3 & 11 \\ 9 & -15 \end{bmatrix} = (AB)'$

Hence proved.

(iii) We have to verify that, $(kA)' = (kA)'$

$$\text{Now, } (kA) = \begin{bmatrix} 0 & -k & 2k \\ 4k & 3k & -4k \end{bmatrix}$$

$$\text{And, } (kA)' = \begin{bmatrix} 0 & 4k \\ -k & 3k \\ 2k & -4k \end{bmatrix}$$

$$\text{Also, } kA' = \begin{bmatrix} 0 & 4k \\ -k & 3k \\ 2k & -4k \end{bmatrix} = (kA)'$$

Hence proved.

28. If $A = \begin{bmatrix} 1 & 2 \\ 4 & 1 \\ 5 & 6 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 2 \\ 6 & 4 \\ 7 & 3 \end{bmatrix}$, then verify that:

(i) $(2A + B)' = 2A' + B'$

(ii) $(A - B)' = A' - B'$

Solution:

$$\text{Given, } A = \begin{bmatrix} 1 & 2 \\ 4 & 1 \\ 5 & 6 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 2 \\ 6 & 4 \\ 7 & 3 \end{bmatrix}$$

$$(i) (2A + B) = \begin{bmatrix} 2 & 4 \\ 8 & 2 \\ 10 & 12 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 6 & 4 \\ 7 & 3 \end{bmatrix} = \begin{bmatrix} 3 & 6 \\ 14 & 6 \\ 17 & 15 \end{bmatrix}$$

$$\text{And, } (2A + B)' = \begin{bmatrix} 3 & 14 & 17 \\ 6 & 6 & 15 \end{bmatrix}$$

$$\text{Also, } 2A' + B'$$

$$= \begin{bmatrix} 2 & 8 & 10 \\ 4 & 2 & 12 \end{bmatrix} + \begin{bmatrix} 1 & 6 & 7 \\ 2 & 4 & 3 \end{bmatrix} = \begin{bmatrix} 3 & 14 & 17 \\ 6 & 6 & 15 \end{bmatrix} = (2A + B)'$$

$$\text{Hence, } 2A' + B' = (2A + B)'$$

$$(ii) \quad A - B = \begin{bmatrix} 1 & 2 \\ 4 & 1 \\ 5 & 6 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ 6 & 4 \\ 7 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ -2 & -3 \\ -2 & 3 \end{bmatrix}$$

$$\text{And, } (A - B)' = \begin{bmatrix} 0 & -2 & -2 \\ 0 & -3 & 3 \end{bmatrix}$$

$$\text{Also, } A' - B' = \begin{bmatrix} 1 & 4 & 5 \\ 2 & 1 & 6 \end{bmatrix} - \begin{bmatrix} 1 & 6 & 7 \\ 2 & 4 & 3 \end{bmatrix} = \begin{bmatrix} 0 & -2 & -2 \\ 0 & -3 & 3 \end{bmatrix} = (A - B)'$$

$$\text{Thus, } A' - B' = (A - B)'$$

- Hence proved

29. Show that $A'A$ and AA' are both symmetric matrices for any matrix A .

Solution:

Let $P = A'A$

So, $P' = (A'A)'$

$$= A'(A')' \quad [\text{As } (AB)' = B'A']$$

Hence, $A'A$ is symmetric matrix for any matrix A .

Now, let $Q = AA'$

So, $Q' = (AA')' = (A')' = AA' = Q$

Hence, AA' is symmetric matrix for any matrix A .

30. Let A and B be square matrices of the order 3×3 . Is $(AB)^2 = A^2 B^2$? Give reasons.

Solution:

As, A and B be square matrices of order 3×3 .

We have, $(AB)^2 = AB \cdot AB$

$$= A(BA)B$$

$$= A(AB)B \quad [\text{If } AB = BA]$$

$$= AAB B$$

$$= A^2 B^2$$

Thus, $(AB)^2 = A^2 B^2$ is true only if $AB = BA$.

31. Show that if A and B are square matrices such that $AB = BA$, then

$$(A + B)^2 = A^2 + 2AB + B^2.$$

Solution:

Given, A and B are square matrices such that $AB = BA$.

So, $(A + B)^2 = (A + B) \cdot (A + B)$

$$= A^2 + AB + BA + B^2$$

$$= A^2 + AB + AB + B^2$$

[Since, $AB = BA$]

$$= A^2 + 2AB + B^2$$

32. Let $A = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 0 \\ 1 & 5 \end{bmatrix}$, $C = \begin{bmatrix} 2 & 0 \\ 1 & -2 \end{bmatrix}$ and $a = 4, b = -2$.

Show that:

(a) $A + (B + C) = (A + B) + C$

(b) $A(BC) = (AB)C$

(c) $(a + b)B = aB + bB$

(d) $a(C - A) = aC - aA$

(e) $(A^T)^T = A$

(f) $(bA)^T = bA^T$

(g) $(AB)^T = B^T A^T$

(h) $(A - B)C = AC - BC$

(i) $(A - B)^T = A^T - B^T$

Solution:

(a) $A + (B + C) = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix} + \begin{bmatrix} 6 & 0 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 7 & 2 \\ 1 & 6 \end{bmatrix}$

And,

$(A + B) + C = \begin{bmatrix} 5 & 2 \\ 0 & 8 \end{bmatrix} + \begin{bmatrix} 2 & 0 \\ 1 & -2 \end{bmatrix} = \begin{bmatrix} 7 & 2 \\ 1 & 6 \end{bmatrix} = A + (B + C)$

(b) $(BC) = \begin{bmatrix} 4 & 0 \\ 1 & 5 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 1 & -2 \end{bmatrix} = \begin{bmatrix} 8 & 0 \\ 7 & -10 \end{bmatrix}$

And, $A(BC) = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 8 & 0 \\ 7 & -10 \end{bmatrix} = \begin{bmatrix} 8 + 14 & 0 - 20 \\ -8 + 21 & 0 - 30 \end{bmatrix} = \begin{bmatrix} 22 & -20 \\ 13 & -30 \end{bmatrix}$

Also, $AB = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 1 & 5 \end{bmatrix} = \begin{bmatrix} 4 + 2 & 0 + 10 \\ -4 + 3 & 0 + 15 \end{bmatrix} = \begin{bmatrix} 6 & 10 \\ -1 & 15 \end{bmatrix}$

Thus, $(AB)C = \begin{bmatrix} 6 & 10 \\ -1 & 15 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 1 & -2 \end{bmatrix} = \begin{bmatrix} 22 & -20 \\ 13 & -30 \end{bmatrix} = A(BC)$

Hence proved.

(c) $(a + b)B = (4 - 2) \begin{bmatrix} 4 & 0 \\ 1 & 5 \end{bmatrix} \quad [\because \text{given } a = 4, b = -2]$
 $= \begin{bmatrix} 8 & 0 \\ 2 & 10 \end{bmatrix}$

Also,

$aB + bB = 4B - 2B = \begin{bmatrix} 16 & 0 \\ 4 & 20 \end{bmatrix} - \begin{bmatrix} 8 & 0 \\ 2 & 10 \end{bmatrix} = \begin{bmatrix} 8 & 0 \\ 2 & 10 \end{bmatrix} = (a + b)B$

Hence proved.

$$(d) \quad C - A = \begin{bmatrix} 2 & 0 \\ 1 & -2 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix} = \begin{bmatrix} 1 & -2 \\ 2 & -5 \end{bmatrix}$$

$$\text{And, } a(C - A) = 4(C - A)$$

$$= \begin{bmatrix} 4 & -8 \\ 8 & -20 \end{bmatrix}$$

$$\text{Also, } aC - aA = 4C - 4A$$

$$= \begin{bmatrix} 8 & 0 \\ 4 & -8 \end{bmatrix} - \begin{bmatrix} 4 & 8 \\ -4 & 12 \end{bmatrix} = \begin{bmatrix} 4 & -8 \\ 8 & -20 \end{bmatrix} = a(C - A)$$

Hence proved.

$$(e) \quad A^T = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}^T = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$$

$$\text{Thus, } (A^T)^T = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}^T = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix} = A$$

Hence proved.

$$(f) \quad (bA)^T = \begin{bmatrix} -2 & -4 \\ 2 & -6 \end{bmatrix}^T \quad [\because b = -2]$$

$$= \begin{bmatrix} -2 & 2 \\ -4 & -6 \end{bmatrix}$$

$$\text{And, } A^T = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$$

$$\text{Thus, } bA^T = \begin{bmatrix} -2 & 2 \\ -4 & -6 \end{bmatrix} = (bA)^T$$

Hence proved.

$$(g) \quad AB = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 1 & 5 \end{bmatrix} = \begin{bmatrix} 4+2 & 0+10 \\ -4+3 & 0+15 \end{bmatrix} = \begin{bmatrix} 6 & 10 \\ -1 & 15 \end{bmatrix}$$

$$\text{So, } (AB)^T = \begin{bmatrix} 6 & -1 \\ 10 & 15 \end{bmatrix}$$

$$\text{Now, } B^T A^T = \begin{bmatrix} 4 & 1 \\ 0 & 5 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 4+2 & -4+3 \\ 0+10 & 0+15 \end{bmatrix} = \begin{bmatrix} 6 & -1 \\ 10 & 15 \end{bmatrix} = (AB)^T$$

Hence proved.

$$(h) (A - B) = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix} - \begin{bmatrix} 4 & 0 \\ 1 & 5 \end{bmatrix} = \begin{bmatrix} 1-4 & 2-0 \\ -1-1 & 3-5 \end{bmatrix} = \begin{bmatrix} -3 & 2 \\ -2 & -2 \end{bmatrix}$$

$$\text{Thus, } (A - B)C = \begin{bmatrix} -3 & 2 \\ -2 & -2 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 1 & -2 \end{bmatrix} = \begin{bmatrix} -4 & -4 \\ -6 & 4 \end{bmatrix}$$

Now,

$$AC = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 1 & -2 \end{bmatrix} = \begin{bmatrix} 4 & -4 \\ 1 & -6 \end{bmatrix}$$

And,

$$BC = \begin{bmatrix} 4 & 0 \\ 1 & 5 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 1 & -2 \end{bmatrix} = \begin{bmatrix} 8 & 0 \\ 7 & -10 \end{bmatrix}$$

Therefore,

$$AC - BC = \begin{bmatrix} 4-8 & -4-0 \\ 1-7 & -6+10 \end{bmatrix} = \begin{bmatrix} -4 & -4 \\ -6 & 4 \end{bmatrix} = (A - B)C$$

Hence proved.

$$(i) (A - B)^T = \begin{bmatrix} 1-4 & 2-0 \\ -1-1 & 3-5 \end{bmatrix}^T = \begin{bmatrix} -3 & 2 \\ -2 & -2 \end{bmatrix}^T = \begin{bmatrix} -3 & -2 \\ 2 & -2 \end{bmatrix}$$

$$\text{Now, } A^T - B^T = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} - \begin{bmatrix} 4 & 1 \\ 0 & 5 \end{bmatrix} = \begin{bmatrix} -3 & -2 \\ 2 & -2 \end{bmatrix} = (A - B)^T$$

Hence proved.

$$33. \text{ If } A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}, \text{ then show that } A^2 = \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}.$$

Solution:

$$\text{Given, } A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

$$\text{Now, } A^2 = A \cdot A$$

$$= \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 \theta - \sin^2 \theta & \cos \theta \sin \theta + \sin \theta \cos \theta \\ -\sin \theta \cos \theta - \cos \theta \sin \theta & -\sin^2 \theta + \cos^2 \theta \end{bmatrix}$$

$$= \begin{bmatrix} \cos 2\theta & 2 \sin \theta \cos \theta \\ 2 \sin \theta \cos \theta & \cos 2\theta \end{bmatrix} = \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$$

Hence proved.

34. If $A = \begin{bmatrix} 0 & -x \\ x & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ and $x^2 = -1$, then show that $(A + B)^2 = A^2 + B^2$

Solution:

Given, $A = \begin{bmatrix} 0 & -x \\ x & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ and $x^2 = -1$

So, $(A + B) = \begin{bmatrix} 0 & -x+1 \\ x+1 & 0 \end{bmatrix}$

And, $(A + B)^2 = \begin{bmatrix} 0 & -x+1 \\ x+1 & 0 \end{bmatrix} \begin{bmatrix} 0 & -x+1 \\ x+1 & 0 \end{bmatrix} = \begin{bmatrix} 1-x^2 & 0 \\ 0 & 1-x^2 \end{bmatrix} \dots(i)$

Also, $A^2 = A \cdot A = \begin{bmatrix} 0 & -x \\ x & 0 \end{bmatrix} \begin{bmatrix} 0 & -x \\ x & 0 \end{bmatrix} = \begin{bmatrix} -x^2 & 0 \\ 0 & -x^2 \end{bmatrix}$

And, $B^2 = B \cdot B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

Thus, $A^2 + B^2 = \begin{bmatrix} -x^2 & 0 \\ 0 & -x^2 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1-x^2 & 0 \\ 0 & 1-x^2 \end{bmatrix} \dots(ii)$

Hence, from equations (i) and (ii), we have

$$(A + B)^2 = A^2 + B^2$$

35. Verify that $A^2 = I$ when $A = \begin{bmatrix} 0 & 1 & -1 \\ 4 & -3 & 4 \\ 3 & -3 & 4 \end{bmatrix}$.

Solution:

Given, $A = \begin{bmatrix} 0 & 1 & -1 \\ 4 & -3 & 4 \\ 3 & -3 & 4 \end{bmatrix}$

So, $A^2 = \begin{bmatrix} 0 & 1 & -1 \\ 4 & -3 & 4 \\ 3 & -3 & 4 \end{bmatrix} \cdot \begin{bmatrix} 0 & 1 & -1 \\ 4 & -3 & 4 \\ 3 & -3 & 4 \end{bmatrix}$

$$= \begin{bmatrix} 0+4-3 & 0-3+3 & 0+4-4 \\ 0-12+12 & 4+9-12 & -4-12+16 \\ 0-12+12 & 3+9-12 & -3-12+16 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

Hence proved.

36. Prove by Mathematical Induction that $(A')^n = (A^n)'$, where $n \in \mathbb{N}$ for any square matrix A .

Solution:

Let $P(n): (A')^n = (A^n)'$

So, $P(1): (A') = (A)'$

$$A' = A'$$

Hence, $P(1)$ is true.

Now, let $P(k) = (A')^k = (A^k)'$, where $k \in \mathbb{N}$

And,

$$\begin{aligned} P(k+1): (A')^{k+1} &= (A')^k A' \\ &= (A^k)' A' \\ &= (AA^k)' \\ &= (A^{k+1})' \end{aligned}$$

Hence, $P(1)$ is true and whenever $P(k)$ is true $P(k+1)$ is true.

Therefore, $P(n)$ is true for all $n \in \mathbb{N}$.

37. Find inverse, by elementary row operations (if possible), of the following matrices

(i) $\begin{bmatrix} 1 & 3 \\ -5 & 7 \end{bmatrix}$ (ii) $\begin{bmatrix} 1 & -3 \\ -2 & 6 \end{bmatrix}$

Solution:

(i) Let $A = \begin{bmatrix} 1 & 3 \\ -5 & 7 \end{bmatrix}$

Now,

$$\begin{bmatrix} 1 & 3 \\ -5 & 7 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} A$$

$$\begin{bmatrix} 1 & 3 \\ 0 & 22 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix} A \quad [\because R_2 \rightarrow R_2 + 5R_1]$$

$$\begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 5/22 & 1/22 \end{bmatrix} A \quad \left[\because R_2 \rightarrow \frac{1}{22} R_2 \right]$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 7/22 & -3/22 \\ 5/22 & 1/22 \end{bmatrix} A \quad [\because R_1 \rightarrow R_1 - 3R_2]$$

$I = BA$, where B is the inverse of A .

Hence,

$$B = \frac{1}{22} \begin{bmatrix} 7 & -3 \\ 5 & -1 \end{bmatrix}$$

(ii) Let $A = \begin{bmatrix} 1 & -3 \\ -2 & 6 \end{bmatrix}$

Now, $\begin{bmatrix} 1 & -3 \\ -2 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} A$

$$\begin{bmatrix} 1 & -3 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} A$$

$$[\because R_2 \rightarrow R_2 + 2R_1]$$

As we obtain all the zeroes in a row of the matrix A on the L.H.S., A^{-1} does not exist.

38. If $\begin{bmatrix} xy & 4 \\ z+6 & x+y \end{bmatrix} = \begin{bmatrix} 8 & w \\ 0 & 6 \end{bmatrix}$, then find values of x, y, z and w.

Solution:

In the given matrix equation,

Comparing the corresponding elements, we get

$$x + y = 6,$$

$$xy = 8,$$

$$z + 6 = 0 \text{ and}$$

$$w = 4$$

From the first two equations, we have

$$(6 - y) \cdot y = 8$$

$$y^2 - 6y + 8 = 0$$

$$(y - 2)(y - 4) = 0$$

$$y = 2 \text{ or } y = 4$$

$$\text{Hence, } x = 4 \text{ and } x = 2$$

$$\text{Also, } z + 6 = 0$$

$$z = -6 \text{ and } w = 4$$

Therefore,

$$x = 2, y = 4 \text{ or } x = 4, y = 2, z = -6 \text{ and } w = 4$$

39. If $A = \begin{bmatrix} 1 & 5 \\ 7 & 12 \end{bmatrix}$ and $B = \begin{bmatrix} 9 & 1 \\ 7 & 8 \end{bmatrix}$, find a matrix C such that $3A + 5B + 2C$ is a null matrix.

Solution:

Let's consider a matrix C, such that

$$C = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$3A + 5B + 2C = O$$

$$\begin{bmatrix} 3 & 15 \\ 21 & 36 \end{bmatrix} + \begin{bmatrix} 45 & 5 \\ 35 & 40 \end{bmatrix} + \begin{bmatrix} 2a & 2b \\ 2c & 2d \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 48 + 2a & 20 + 2b \\ 56 + 2c & 76 + 2d \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Comparing the terms,

$$2a + 48 = 0 \Rightarrow a = -24$$

$$20 + 2b = 0 \Rightarrow b = -10$$

$$56 + 2c = 0 \Rightarrow c = -28$$

And,

$$76 + 2d = 0 \Rightarrow d = -38$$

Therefore, the matrix C is

$$C = \begin{bmatrix} -24 & -10 \\ -28 & -38 \end{bmatrix}$$

40. If $A = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$, then find $A^2 - 5A - 14I$. Hence, obtain A^3 .

Solution:

Given, $A = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$... (i)

Now, $A^2 = A \cdot A = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix} \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix} = \begin{bmatrix} 29 & -25 \\ -20 & 24 \end{bmatrix}$

So, $A^2 - 5A - 14I = \begin{bmatrix} 29 & -25 \\ -20 & 24 \end{bmatrix} - \begin{bmatrix} 15 & -25 \\ -20 & 10 \end{bmatrix} - \begin{bmatrix} 14 & 0 \\ 0 & 14 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

$$A \cdot A^2 - 5A \cdot A = 14AI = O$$

$$A^3 - 5A^2 - 14A = O$$

$$A^3 = 5A^2 + 14A$$

$$= 5 \begin{bmatrix} 29 & -25 \\ -20 & 24 \end{bmatrix} + 14 \begin{bmatrix} 3 & -5 \\ -4 & -2 \end{bmatrix}$$

$$= \begin{bmatrix} 145 & -125 \\ -100 & 120 \end{bmatrix} + \begin{bmatrix} 42 & -70 \\ -56 & 28 \end{bmatrix}$$

Hence, $A^3 = \begin{bmatrix} 187 & -195 \\ -156 & 148 \end{bmatrix}$

41. Find the value of a, b, c and d, if

$$3 \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & 6 \\ -1 & 2d \end{bmatrix} + \begin{bmatrix} 4 & a+b \\ c+d & 3 \end{bmatrix}.$$

Solution:

Given,

$$3 \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & 6 \\ -1 & 2d \end{bmatrix} + \begin{bmatrix} 4 & a+b \\ c+d & 3 \end{bmatrix}$$

$$\begin{bmatrix} 3a & 3b \\ 3c & 3d \end{bmatrix} = \begin{bmatrix} a+4 & 6+a+b \\ -1+c+d & 2d+3 \end{bmatrix}$$

Now,

$$3a = a + 4 \Rightarrow a = 2$$

$$3b = 6 + a + b$$

$$3b - b = 8$$

$$\Rightarrow b = 4$$

And,

$$3d = 3 + 2d$$

$$\Rightarrow d = 3$$

And,

$$3c = c + d - 1$$

$$2c = 3 - 1 = 2$$

$$\Rightarrow c = 1$$

Hence,

$$a = 2, b = 4, c = 1 \text{ and } d = 3$$

42. Find the matrix A such that

$$\begin{bmatrix} 2 & -1 \\ 1 & 0 \\ -3 & 4 \end{bmatrix} A = \begin{bmatrix} -1 & -8 & -10 \\ 1 & -2 & -5 \\ 9 & 22 & 15 \end{bmatrix}.$$

Solution:

Let, $A = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix}$

$$\begin{bmatrix} 2 & -1 \\ 1 & 0 \\ -3 & 4 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} = \begin{bmatrix} -1 & -8 & -10 \\ 1 & -2 & -5 \\ 9 & 22 & 15 \end{bmatrix}$$

$$\begin{bmatrix} 2a-d & 2b-e & 2c-f \\ a & b & c \\ -3a+4d & -3b+4e & -3c+4f \end{bmatrix} = \begin{bmatrix} -1 & -8 & -10 \\ 1 & -2 & -5 \\ 9 & 22 & 15 \end{bmatrix}$$

Now, by equality of matrices, we get

$$a = 1, b = -2, c = -5$$

And,

$$2a - d = -1 \Rightarrow d = 2a + 1 = 3;$$

$$2b - e = -8 \Rightarrow e = 2(-2) + 8 = 4$$

$$2c - f = -10 \Rightarrow f = 2c + 10 = 0$$

Thus,

$$A = \begin{bmatrix} 1 & -2 & -5 \\ 3 & 4 & 0 \end{bmatrix}$$

43. If $A = \begin{bmatrix} 1 & 2 \\ 4 & 1 \end{bmatrix}$, find $A^2 + 2A + 7I$.

Solution:

Given,

$$A = \begin{bmatrix} 1 & 2 \\ 4 & 1 \end{bmatrix}$$

$$A^2 = A \cdot A = \begin{bmatrix} 1 & 2 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 4 & 1 \end{bmatrix} = \begin{bmatrix} 1+8 & 2+2 \\ 4+4 & 8+1 \end{bmatrix} = \begin{bmatrix} 9 & 4 \\ 8 & 9 \end{bmatrix}$$

$$A^2 + 2A + 7I = \begin{bmatrix} 9 & 4 \\ 8 & 9 \end{bmatrix} + \begin{bmatrix} 2 & 4 \\ 8 & 2 \end{bmatrix} + \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} = \begin{bmatrix} 18 & 8 \\ 16 & 18 \end{bmatrix}$$

44. If $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$, and $A^{-1} = A'$, find value of α .

Solution:

Given,

$$A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix} \text{ and } A' = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

Also,

$$A^{-1} = A'$$

$$AA^{-1} = AA'$$

$$I = AA'$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \cos^2 \alpha + \sin^2 \alpha & 0 \\ 0 & \sin^2 \alpha + \cos^2 \alpha \end{bmatrix}$$

By using equality of matrices, we get
 $\cos^2 \alpha + \sin^2 \alpha = 1$, which is true for all real values of α .

45. If the matrix $\begin{bmatrix} 0 & a & 3 \\ 2 & b & -1 \\ c & 1 & 0 \end{bmatrix}$ is a skew symmetric matrix, find the values of a , b and c .

Solution:

$$\text{Let } A = \begin{bmatrix} 0 & a & 3 \\ 2 & b & -1 \\ c & 1 & 0 \end{bmatrix}$$

As, A is skew - symmetric matrix.

So, we have

$$A' = -A$$

Then,

$$\Rightarrow \begin{bmatrix} 0 & 2 & c \\ a & b & 1 \\ 3 & -1 & 0 \end{bmatrix} = - \begin{bmatrix} 0 & a & 3 \\ 2 & b & -1 \\ c & 1 & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 0 & 2 & c \\ a & b & 1 \\ 3 & -1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -a & -3 \\ -2 & -b & 1 \\ -c & -1 & 0 \end{bmatrix}$$

By equality of matrices, we get

$$a = -2, c = -3 \text{ and } b = -b \Rightarrow b = 0$$

Hence,

$$a = -2, b = 0 \text{ and } c = -3$$

46. If $P(x) = \begin{bmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{bmatrix}$, then show that

$$P(x) \cdot P(y) = P(x+y) = P(y) \cdot P(x).$$

Solution:

Given,

$$P(x) = \begin{bmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{bmatrix}$$

$$\text{So, } P(y) = \begin{bmatrix} \cos y & \sin y \\ -\sin y & \cos y \end{bmatrix}$$

Now,

$$\begin{aligned}
 P(x) \cdot P(y) &= \begin{bmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{bmatrix} \begin{bmatrix} \cos y & \sin y \\ -\sin y & \cos y \end{bmatrix} \\
 &= \begin{bmatrix} \cos x \cdot \cos y - \sin x \cdot \sin y & \cos x \cdot \sin y + \sin x \cdot \cos y \\ -\sin x \cdot \cos y - \cos x \cdot \sin y & -\sin x \cdot \sin y + \cos x \cdot \cos y \end{bmatrix} \\
 &= \begin{bmatrix} \cos(x+y) & \sin(x+y) \\ -\sin(x+y) & \cos(x+y) \end{bmatrix} \\
 &= P(x+y) \quad \dots(i)
 \end{aligned}$$

Also,

$$\begin{aligned}
 P(y) \cdot P(x) &= \begin{bmatrix} \cos y & \sin y \\ -\sin y & \cos y \end{bmatrix} \begin{bmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{bmatrix} \\
 &= \begin{bmatrix} \cos y \cdot \cos x - \sin y \cdot \sin x & \cos y \cdot \sin x + \sin y \cdot \cos x \\ -\sin y \cdot \cos x - \sin x \cdot \cos y & -\sin y \cdot \sin x + \cos y \cdot \cos x \end{bmatrix} \\
 &= \begin{bmatrix} \cos(x+y) & \sin(x+y) \\ -\sin(x+y) & \cos(x+y) \end{bmatrix} \quad \dots(ii)
 \end{aligned}$$

Therefore, from (i) and (ii), we get

$$P(x) \cdot P(y) = P(x+y) = P(y) \cdot P(x)$$

47. If A is square matrix such that $A^2 = A$, show that $(I + A)^3 = 7A + I$.

Solution:

We know that,

$$A \cdot I = I \cdot A$$

So, A and I are commutative.

Thus, we can expand $(I + A)^3$ like real numbers expansion.

$$\begin{aligned}
 \text{So, } (I + A)^3 &= I^3 + 3I^2A + 3IA^2 + A^3 \\
 &= I + 3IA + 3A^2 + AA^2 \quad (\text{As } I^n = I, n \in \mathbb{N}) \\
 &= I + 3A + 3A + AA \\
 &= I + 3A + 3A + A^2 = I + 3A + 3A + A = I + 7A
 \end{aligned}$$

48. If A, B are square matrices of same order and B is a skew-symmetric matrix, show that $A'BA$ is skew symmetric.

Solution:

Given, A and B are square matrices such that B is a skew-symmetric matrix

$$\text{So, } B' = -B$$

Now, we have to prove that $A'BA$ is a skew-symmetric matrix.

$$\begin{aligned}
 (A'BA)' &= A'B'(A')' && [\text{Since, } (AB)' = B'A'] \\
 &= A'(-B)A \\
 &= -A'BA
 \end{aligned}$$

Hence, $A'BA$ is a skew-symmetric matrix.

Long Answer (L.A)

49. If $AB = BA$ for any two square matrices, prove by mathematical induction that $(AB)^n = A^n B^n$.

Solution:

Let $P(n) : (AB)^n = A^n B^n$

So, $P(1) : (AB)^1 = A^1 B^1$

$$AB = AB$$

So, $P(1)$ is true.

Let $P(n)$ is true for some $k \in \mathbb{N}$

Now,

$$\begin{aligned}
 (AB)^{k+1} &= (AB)^k(AB) && (\text{using (i)}) \\
 &= A^k B^k (AB) \\
 &= A^k B^{k-1} (BA) B \\
 &= A^k B^{k-1} (AB) B && (\text{as given } AB = BA) \\
 &= A^k B^{k-1} AB^2 \\
 &= A^k B^{k-2} (BA) B^2 \\
 &= A^k B^{k-2} ABB^2 \\
 &= A^k B^{k-2} AB^3 \\
 &\dots\dots \\
 &\dots\dots \\
 &= A^{k+1} B^{k+1}
 \end{aligned}$$

Hence, $P(1)$ is true and whenever $P(k)$ is true $P(k+1)$ is true.

Thus, $P(n)$ is true for all $n \in \mathbb{N}$.

50. Find x, y, z if $A = \begin{bmatrix} 0 & 2y & z \\ x & y & -z \\ x & -y & z \end{bmatrix}$ satisfies $A' = A^{-1}$.

Solution:

Matrix A is such that $A' = A^{-1}$

$$AA' = I$$

$$\begin{bmatrix} 0 & 2y & z \\ x & y & -z \\ x & -y & z \end{bmatrix} \begin{bmatrix} 0 & x & x \\ 2y & y & -y \\ z & -z & z \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 4y^2 + z^2 & 2y^2 - z^2 & -2y^2 + z^2 \\ 2y^2 - z^2 & x^2 + y^2 + z^2 & x^2 - y^2 - z^2 \\ -2y^2 + z^2 & x^2 - y^2 + z^2 & x^2 + y^2 + z^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$4y^2 + z^2 = 1$$

$$2y^2 - z^2 = 0$$

$$x^2 + y^2 + z^2 = 1$$

$$x^2 - y^2 - z^2 = 0$$

$$y^2 = 1/6, z^2 = 1/3, x^2 = 1/2$$

So, the roots are:

$$x = \pm \frac{1}{\sqrt{2}}$$

$$y = \pm \frac{1}{\sqrt{6}}$$

And,

$$z = \pm \frac{1}{\sqrt{3}}$$

NAGEEN GROUP
tamso ma jyotirgamaya

CLICK TO DOWNLOAD NCERT EXEMPLAR SOLUTIONS

CLASS 12: MATHEMATICS (ALL CHAPTERS)





JOIN OUR WHATSAPP GROUPS

FOR FREE EDUCATIONAL
RESOURCES





JOIN SCHOOL OF EDUCATORS WHATSAPP GROUPS FOR FREE EDUCATIONAL RESOURCES

We are thrilled to introduce the School of Educators WhatsApp Group, a platform designed exclusively for educators to enhance your teaching & Learning experience and learning outcomes. Here are some of the key benefits you can expect from joining our group:

BENEFITS OF SOE WHATSAPP GROUPS

- **Abundance of Content:** Members gain access to an extensive repository of educational materials tailored to their class level. This includes various formats such as PDFs, Word files, PowerPoint presentations, lesson plans, worksheets, practical tips, viva questions, reference books, smart content, curriculum details, syllabus, marking schemes, exam patterns, and blueprints. This rich assortment of resources enhances teaching and learning experiences.
- **Immediate Doubt Resolution:** The group facilitates quick clarification of doubts. Members can seek assistance by sending messages, and experts promptly respond to queries. This real-time interaction fosters a supportive learning environment where educators and students can exchange knowledge and address concerns effectively.
- **Access to Previous Years' Question Papers and Topper Answers:** The group provides access to previous years' question papers (PYQ) and exemplary answer scripts of toppers. This resource is invaluable for exam preparation, allowing individuals to familiarize themselves with the exam format, gain insights into scoring techniques, and enhance their performance in assessments.

- **Free and Unlimited Resources:** Members enjoy the benefit of accessing an array of educational resources without any cost restrictions. Whether its study materials, teaching aids, or assessment tools, the group offers an abundance of resources tailored to individual needs. This accessibility ensures that educators and students have ample support in their academic endeavors without financial constraints.
- **Instant Access to Educational Content:** SOE WhatsApp groups are a platform where teachers can access a wide range of educational content instantly. This includes study materials, notes, sample papers, reference materials, and relevant links shared by group members and moderators.
- **Timely Updates and Reminders:** SOE WhatsApp groups serve as a source of timely updates and reminders about important dates, exam schedules, syllabus changes, and academic events. Teachers can stay informed and well-prepared for upcoming assessments and activities.
- **Interactive Learning Environment:** Teachers can engage in discussions, ask questions, and seek clarifications within the group, creating an interactive learning environment. This fosters collaboration, peer learning, and knowledge sharing among group members, enhancing understanding and retention of concepts.
- **Access to Expert Guidance:** SOE WhatsApp groups are moderated by subject matter experts, teachers, or experienced educators can benefit from their guidance, expertise, and insights on various academic topics, exam strategies, and study techniques.

Join the School of Educators WhatsApp Group today and unlock a world of resources, support, and collaboration to take your teaching to new heights. To join, simply click on the group links provided below or send a message to +91-95208-77777 expressing your interest.

**Together, let's empower ourselves & Our Students and
inspire the next generation of learners.**

**Best Regards,
Team
School of Educators**

Join School of Educators WhatsApp Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class WhatsApp Group.

Kindergarten to Class XII (For Teachers Only)



[Click Here to Join](#)

Class 1



[Click Here to Join](#)

Class 2



[Click Here to Join](#)

Class 3



[Click Here to Join](#)

Class 4



[Click Here to Join](#)

Class 5



[Click Here to Join](#)

Class 6



[Click Here to Join](#)

Class 7



[Click Here to Join](#)

Class 8



[Click Here to Join](#)

Class 9



[Click Here to Join](#)

Class 10



[Click Here to Join](#)

Class 11 (Science)



[Click Here to Join](#)

Class 11 (Humanities)



[Click Here to Join](#)

Class 11 (Commerce)



[Click Here to Join](#)

Class 12 (Science)



[Click Here to Join](#)

Class 12 (Humanities)



[Click Here to Join](#)

Class 12 (Commerce)



[Click Here to Join](#)

Kindergarten

Subject Wise Secondary and Senior Secondary Groups (IX & X For Teachers Only) Secondary Groups (IX & X)



[Click Here to Join](#)

SST



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Science



[Click Here to Join](#)

English



[Click Here to Join](#)

Hindi-A



[Click Here to Join](#)

IT Code-402



[Click Here to Join](#)

Hindi-B



[Click Here to Join](#)

Artificial Intelligence

Senior Secondary Groups (XI & XII For Teachers Only)



[Click Here to Join](#)

Physics



[Click Here to Join](#)

Chemistry



[Click Here to Join](#)

English



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Biology



[Click Here to Join](#)

Accountancy



[Click Here to Join](#)

Economics



[Click Here to Join](#)

BST



[Click Here to Join](#)

History



[Click Here to Join](#)

Geography



[Click Here to Join](#)

Sociology



[Click Here to Join](#)

Hindi Elective



[Click Here to Join](#)

Hindi Core



[Click Here to Join](#)

Home Science



[Click Here to Join](#)

Sanskrit



[Click Here to Join](#)

Psychology



[Click Here to Join](#)

Political Science



[Click Here to Join](#)

Painting



[Click Here to Join](#)

Vocal Music



[Click Here to Join](#)

Comp. Science



[Click Here to Join](#)

IP



[Click Here to Join](#)

Physical Education



[Click Here to Join](#)

APP. Mathematics



[Click Here to Join](#)

Legal Studies



[Click Here to Join](#)

Entrepreneurship



[Click Here to Join](#)

French



[Click Here to Join](#)

IT



[Click Here to Join](#)

Artificial Intelligence

Other Important Groups (For Teachers & Principal's)



[Click Here to Join](#)

Principal's Group



[Click Here to Join](#)

Teachers Jobs



[Click Here to Join](#)

IIT/NEET

Join School of Educators WhatsApp Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class WhatsApp Group.

Kindergarten to Class XII (For Students Only)



[Click Here to Join](#)

Class 1



[Click Here to Join](#)

Class 2



[Click Here to Join](#)

Class 3



[Click Here to Join](#)

Class 4



[Click Here to Join](#)

Class 5



[Click Here to Join](#)

Class 6



[Click Here to Join](#)

Class 7



[Click Here to Join](#)

Class 8



[Click Here to Join](#)

Class 9



[Click Here to Join](#)

Class 10



[Click Here to Join](#)

Class 11 (Science)



[Click Here to Join](#)

Class 11 (Humanities)



[Click Here to Join](#)

Class 11 (Commerce)



[Click Here to Join](#)

Class 12 (Science)



[Click Here to Join](#)

Class 12 (Humanities)



[Click Here to Join](#)

Class 12 (Commerce)



[Click Here to Join](#)

**Artificial Intelligence
(VI TO VIII)**

Subject Wise Secondary and Senior Secondary Groups (IX & X For Students Only) Secondary Groups (IX & X)



[Click Here to Join](#)

SST



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Science



[Click Here to Join](#)

English



[Click Here to Join](#)

Hindi



[Click Here to Join](#)

IT Code



[Click Here to Join](#)

Artificial Intelligence

Senior Secondary Groups (XI & XII For Students Only)



[Click Here to Join](#)

Physics



[Click Here to Join](#)

Chemistry



[Click Here to Join](#)

English



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Biology



[Click Here to Join](#)

Accountancy



[Click Here to Join](#)

Economics



[Click Here to Join](#)

BST



[Click Here to Join](#)

History



[Click Here to Join](#)

Geography



[Click Here to Join](#)

Sociology



[Click Here to Join](#)

Hindi Elective



[Click Here to Join](#)

Hindi Core



[Click Here to Join](#)

Home Science



[Click Here to Join](#)

Sanskrit



[Click Here to Join](#)

Psychology



[Click Here to Join](#)

Political Science



[Click Here to Join](#)

Painting



[Click Here to Join](#)

Music



[Click Here to Join](#)

Comp. Science



[Click Here to Join](#)

IP



[Click Here to Join](#)

Physical Education



[Click Here to Join](#)

APP. Mathematics



[Click Here to Join](#)

Legal Studies



[Click Here to Join](#)

Entrepreneurship



[Click Here to Join](#)

French



[Click Here to Join](#)

IT



[Click Here to Join](#)

AI



[Click Here to Join](#)

IIT/NEET



[Click Here to Join](#)

CUET

Groups Rules & Regulations:

To maximize the benefits of these WhatsApp groups, follow these guidelines:

1. Share your valuable resources with the group.
2. Help your fellow educators by answering their queries.
3. Watch and engage with shared videos in the group.
4. Distribute WhatsApp group resources among your students.
5. Encourage your colleagues to join these groups.

Additional notes:

1. Avoid posting messages between 9 PM and 7 AM.
2. After sharing resources with students, consider deleting outdated data if necessary.
3. It's a NO Nuisance groups, single nuisance and you will be removed.
 - No introductions.
 - No greetings or wish messages.
 - No personal chats or messages.
 - No spam. Or voice calls
 - Share and seek learning resources only.

Please only share and request learning resources. For assistance, contact the helpline via WhatsApp: +91-95208-77777.

Join Premium WhatsApp Groups Ultimate Educational Resources!!

Join our premium groups and just Rs. 1000 and gain access to all our exclusive materials for the entire academic year. Whether you're a student in Class IX, X, XI, or XII, or a teacher for these grades, Artham Resources provides the ultimate tools to enhance learning. Pay now to delve into a world of premium educational content!

[Click here for more details](#)



Click Here to Join

Class 9



Click Here to Join

Class 10



Click Here to Join

Class 11



Click Here to Join

Class 12

📢 Don't Miss Out! Elevate your academic journey with top-notch study materials and secure your path to top scores! Revolutionize your study routine and reach your academic goals with our comprehensive resources. Join now and set yourself up for success! 🇧🇩🌟

Best Wishes,

Team

School of Educators & Artham Resources

SKILL MODULES BEING OFFERED IN MIDDLE SCHOOL



Artificial Intelligence



Beauty & Wellness




Design Thinking & Innovation



Financial Literacy



Handicrafts



Information Technology



Marketing/Commercial Application




Mass Media - Being Media Literate




Travel & Tourism




Coding



Data Science (Class VIII only)




Augmented Reality / Virtual Reality



Digital Citizenship




Life Cycle of Medicine & Vaccine



Things you should know about keeping Medicines at home




What to do when Doctor is not around



Humanity & Covid-19



Blue Pottery




Pottery



Block Printing




Food



Food Preservation



Baking



Herbal Heritage



Khadi




Mask Making




Mass Media



Making of a Graphic Novel



Kashmiri Embroidery




Embroidery




Rockets



Satellites




Application of Satellites




Photography

SKILL SUBJECTS AT SECONDARY LEVEL (CLASSES IX – X)



Retail



Information Technology




Security




Automotive




Introduction To Financial Markets



Introduction To Tourism



Beauty & Wellness



Agriculture



Food Production



Front Office Operations



Banking & Insurance



Marketing & Sales



Health Care



Apparel



Multi Media



Multi Skill Foundation Course



Artificial Intelligence



Physical Activity Trainer



Data Science



Electronics & Hardware (NEW)




Foundation Skills For Sciences (Pharmaceutical & Biotechnology)(NEW)




Design Thinking & Innovation (NEW)

SKILL SUBJECTS AT SR. SEC. LEVEL (CLASSES XI – XII)



Retail



Information Technology



Web Application



Automotive



Financial Markets Management



Tourism



Beauty & Wellness



Agriculture



Food Production



Front Office Operations



Banking



Marketing




Health Care




Insurance




Horticulture



Typography & Comp.
Application




Geospatial Technology



Electrical Technology



Electronic Technology



Multi-Media




Taxation



Cost Accounting



Office Procedures & Practices



Shorthand (English)




Shorthand (Hindi)



Air-Conditioning & Refrigeration



Medical Diagnostics



Textile Design




Design



Salesmanship



Business Administration



Food Nutrition & Dietetics



Mass Media Studies



Library & Information Science



Fashion Studies



Applied Mathematics



Yoga



Early Childhood Care & Education




Artificial Intelligence



Data Science



Physical Activity Trainer(new)





Land Transportation Associate (NEW)



Electronics & Hardware (NEW)



Design Thinking & Innovation (NEW)



**Join our
Telegram Group
for more sample
papers**



**Join our
Telegram Channel
for more books &
Notes**

- Pdf Purchase

- Print Purchase

- Amazon Print Purchase



Scan QR Code to join
Telegram Groups &
Channels for more free
resources.

